Biology of Sexual Development

Objectives:

- 1. Sex vs. Gender defined.
- 2. Biological sex based on inheritance of sex chromosomes from gametes.
- 3. Biological sex during embryonic development driven by testosterone & estrogen exposure.
- 4. Intersex: exception when biological sex is NOT binary.
- 5. Biological causes of intersex condition.











Reading Assignments:


- 1. Rediscovering Biology: Sex & Gender (Pgs 5 9)
- 2. <u>Gray Areas of Gender Intersex Essay</u>
 (this is a bit edgy for some readers, but important)
- 3. Should We "Fix" Intersex Children?
- 4. Gender Definitions article
- 5. Hormone Hell
- 6. Gender-Bender Chemicals Are Harming Unborn Boys
- 7. Boys Won't Be Boys
























Rediscovering Biology: Sex & Gender Pgs 1, 5 – 9 (topics we are covering are in yellow)



Review:

- Gametes (sex cells) versus somatic cells
- · haploid versus diploid
- role of mitosis & meiosis
- Sexual reproduction
- · Sexual differentiation at 8 weeks:
 - > development from undifferentiated gonad (genital tubercle, labial folds, urethral folds, wolffian and mullerian ducts)
 - >role of SRY & TDF genes
 - > influence of testosterone
 - >influence of estrogen

2. When Sexual Development Goes ... Differently


(or disorders of sexual development, "DSD") =

chromosomal sex is inconsistent with phenotypic sex, or in which the phenotype is not classifiable as either male or female. There is a discrepancy between the external genitals and the internal structures. Occurs in 0.018% of the population

Criteria for being intersex quite diverse, but includes:

- > Having both ovary and testis (internally)
- > External genitals don't match clear male / female
- > Sex chromosomes don't match clear XX or XY.

Source: Intersex Society of North America – www.isna.org/faq



2. When Sexual Development Goes ... Differently

The following is a list of disorders of sex development that sometimes involve intersex anatomy.

- · How do I know if I have an intersex condition?
- ★ 5-alpha reductase deficiency
- Androgen Insensitivity Syndrome (AIS)
 - Aphallia
- Clitoromegaly (large clitoris)
- Congenital Adrenal Hyperplasia (CAH)
 - · gonadal dysgenesis (partial & complete)
- hypospadias
 - . I have a line along the underside of my penis
- * Klinefelter Syndrome
- micropenis
 - · mosaicism involving "sex" chromosomes
 - MRKH (Mullerian agenesis; vaginal agenesis; congenital absence of vagina)
 - · ovo-testes (formerly called "true hermaphroditism")
- Partial Androgen Insensitivity Syndrome (PAIS)
 - Progestin Induced Virilization
 - Swyer Syndrome
- Turner Syndrome

Source: Intersex Society of North America - www.isna.org/faq

Intersex: exception when biological sex is NOT binary.







Micropenis

Hypospadia

Clitoromegaly, hypospadia

Micropenis = _

(1.5 / 10,000 male births)

Cliteromegaly = _

(rare, usually due to hormones (exposure to testosterone) in girls, such as those with congenital adrenal hyperplasia or polycystic ovarian syndrome)

Hypospadia =

(fairly common, 1 / 250 male births)

Possible Causes of Intersex Condition:




- 1. Nondisjunction of sex chromosomes (during meiosis) Meiosis problem.
- 2. Kallman syndrome (hypogonadism) Brain problem.
- 3. Androgen-insensitivity syndrome (AIS) Tissue problem.
- 4. Congenital adrenal hyperplasia (CAH) Adrenal gland problem.
- **5. 5-alpha reductase deficiency** Enzyme problem.
- 6. Endocrine-Disrupting Chemicals Environmental problem

1. Nondisju	ınction i	in the sex	chromosomes
-------------	-----------	------------	-------------

=_____

- > Egg or sperm has extra, or too few, sex chromosomes.
- > May affect as many as 18 % of all ova and 4% of all sperm!

Source: Science Direct



1. Nondisjunction in the sex chromosomes

Example:

A) _____ = XXY male (extra X chromosome)


- > Affects about 1 / 650 newborn boys.
- > It is among the most common sex chromosome disorders.
- > Have smaller testes, & low testosterone.
- > Some degree of feminized features after puberty <u>if not diagnosed & treated</u> with supplemental testosterone.


May or may not also have:

- > Gynocomastia = _____
- > Cryptorchidism = _____
- > Hypospadia =
- > Micropenis = ____
- > Hip development due to estrogen changing fat deposition.z
- > Infertility
- > Taller than average
- > Poor muscle development

Source: https://medlineplus.gov/genetics/condition/klinefelter-syndrome/

Frontal baldness Klinefelter Syndrome absent -Poor beard growth Tendency to grow fewer -Narrow chest hairs shoulders Breastdevelopment Wide Female-type -hips pubic hair pattern-Small testicular Long arms size and legs







= Brain problem - hypothalamus doesn't stimulate gonads.

> Rare - 1 out of 50,000 births

Treatment:

Hormone replacement therapy





4


= fetus with overgrown adrenal cortex. Produces high testosterone. In genetic female (XX) — testosterone masculinizes fetus. Born intersex with ambiguous genitals.



- > 1 / 13,000 girl births
- > Megaloclitoris (enlarged "phalli
- > Full or partial fusion of labia into
- > Studies have shown CAH girls have masculinized brain.




Increased androgen production results in ambiguous genitalia in newborn girls.



6. Endocrine-Disrupting Chemicals (EDCs)

= Man-made chemicals released into environment that mimic or disrupt natural hormones that influence sex development and functions.



6. Endocrine-Disrupting Chemicals (EDCs)

Ex. **Bisphenol** = estrogen-mimic found in plastics. Can feminize males at 30 ug/kg/day. At 20 ug/kg/day can damage eggs & chromosomes. At 2 ug/kg/day can cause early puberty in girls, increase growth prostate and breast cells.



Dose (µg/kg/day)	Effects (measured in studies of laboratory animals)	
0.025	Permanent changes to genital tract	2005
0.025	Changes in breast tissue that predispose cells to hormones and cercinogens	2005
1.6	Low levels of human exposure from diet	2003
2	30% increase in prostate weight	1997
2.4	Signs of early puberty	2002
2.4	Decline in testicular testosterone	2004
2.5	Breast cells predisposed to cancer	2006
10	Prostate cells more sensitive to hormones and cancer	2006
10	Insulin resistance	2006
10	Decreased maternal behavior	2002
13	High levels of human exposure from diet	2003
20	Damage to eggs and chromosomes	2003
25	Health Canada provisional human exposure limit.	1999
30	Hyperactivity	2004
30	Reversal of normal sex difference in brain structure	2001
50	U.S. human exposure limit	1996

Review

5 ways sexual development can go wrong

- 1. Problem in meiosis Non-disjunction of sex chromosomes in eggs or sperm
 - Klinefelter & Turner Syndromes
- 2. Brain problem Kallmann Syndrome
- 3. Problem with tissues AIS
- 4. Problem with adrenal glands CAH
- 5. Enzyme problem $5-\alpha$ reductase deficiency
- 6. Endocrine-Disrupting Chemicals

33