CH. 15 - REPRODUCTIVE SYSTEM

Objectives:


- 1. Review male & female reproductive anatomy
- 2. Gametogenesis & steroidogenesis
- 3. Reproductive problems









1







Epididymis = where sperm mature before leaving in ejaculation (sperm warehouse).

Scrotum = where testes housed outside of abdominal cavity (to keep ~3 degrees

cooler.

Cremaster muscle =

Muscle that lifts or lowers testes to maintain temperature.


Spermatic cord =

Connective tissue that wraps aroune cremaster, testes, & testicular nerve and blood vessels

Inguinal ring =

Opening in inguinal ligament through which testes descend (around 7 months gestation).

Testiculi What is Cryptorchidism? = when 1 or both testes Testiculi retained in the abdomen. 3 – 5% births) Ureter Inguinal ligament Urinary bladder **Inguinal ring** Ductus inguinal ring Structures -Spermatic cord within spermatic Testicular artery-Testicular nerve External spermatic fascia **Cremaster muscle** epididymis Internal spermatic fascia Dartos muscle scrotum



Testicle descends

into scrotum at

about 4-6 weeks

before birth

to top of groin at

about 12 weeks

after conception


6

near the kidney about

7 weeks after conception

3 Sperm Transport Tubes:

- 1. Vas deferens = transports sperm from epididymis to ejaculatory duct.
- **2. Ejaculatory duct =** tube through prostate by which sperm reach the urethra.
- 3. Urethra = common passage for urine and semen (but not at same time)




7


3 Male Secretory Glands:

1. Seminal vesicles = large, paired glands that meet with vas deferens and contribute secretions to seminal fluid.

produces:


- alkaline mucus buffers the sperm from vaginal acidity (Lactobacillus)
- Fructose simple sugar to provide energy to swimming sperm
- Prostaglandin hormone that causes mild uterine contractions (aid sperm transport?)
- Prostate = gland under bladder which secretes mucus.
- **3. Bulbourethral gland =** Gland that secretes lubricating Fluid, to lubricate head of penis, during sexual arousal.





CLINICAL APPLICATIONS


Prostate cancer is commonly tested using a blood test for *prostate-specific antigen (PSA)*. A more common disorder, affecting most men over 60 to different degrees, is **benign prostatic hyperplasia** (**BPH**). This is responsible for most cases of bladder outlet obstruction, causing difficulty in urination. BPH treatment may involve a surgical procedure called *transurethral resection (TUR)*, or the use of drugs. These drugs include α_1 -adrenergic receptor blockers (chapter 6), which decrease the muscle tone of the prostate and bladder neck, making urination easier, and 5α -reductase inhibitors. The latter drugs block the conversion of testosterone into dihydrotestosterone (DHT), which reduces androgen stimulation and thus the size of the prostate.²



How an erection works:

Review of Ch 4 part 1, and Ch 14

- **1. Arousal** Causes **nitric oxide** (**NO**) release in arteries of corpus cavernosa.
- **2.** NO causes production of a chemical messenger called cGMP).
- 3. **cGMP** causes arteries to relax & they open wide (**vasodilate**) allowing blood into spongy chambers.
- **4.** Fluid pressure of blood causes erection.
- 5. When stimulation done, or after ejaculation, cGMP is broken down by enzyme (phosphodiesterase). Erection ends



CLINICAL APPLICATIONS

Nitric oxide, released in the penis in response to parasympathetic nerve activation, enters the smooth muscle cells in the arterioles and stimulates the production of a second messenger, cyclic guanosine monophosphate (cGMP). The cGMP causes the smooth muscle cells to relax and the vessels to dilate, so that more blood can flow to the corpora cavernosa and produce erection. A particular cGMP phosphodiesterase enzyme then breaks down cGMP, ending the erection. Erectile dysfunction is now often treated with drugs such as sildenafil (Viagra), which block the cGMP phosphodiesterase enzyme. These drugs increase the cellular concentration of cGMP and thereby promote erection.

13

How ED Drugs work (Viagra, Cialis, Levitra):

<u>Phosphodiesterase inhibitor</u> = a chemical that inhibits phosphodiesterase.

So ..., what would giving a phosphodiesterase inhibitor do to cGMP levels in the corpus cavernosa?

What would that do to arteries in the penis? Vasadlated

What would that do w/respect to an erection? helps one

Viagra, Cialis, & Levitra are phosphodiesterase inhibitors.

Side effects: drop in blood pressure (dizziness or loss of vision) AND Priapism = Erection longer than 3-5 hrs

Click **HERE** to read more about priapism (causes, repair, etc...)

4. Gametogenesis and Steroidogenesis in Males and Females

Gametogenesis = making (maturing) of eggs and sperm.

- > Spermatogenesis = maturing sperm
- > Oogenesis = maturing eggs


Steroidogenesis = making steroid hormones (testosterone, estrogen, and progesterone).


Hypothalamus (ventromedial nucleus) > GnRH > pituitary > LH & FSH


LH – causes testosterone production in testes Leydig cells, estrogen production in ovaries & ovulation.


FSH – causes egg and sperm maturation


15














CLINICAL APPLICATIONS

About 60 million women worldwide currently use oral contraceptives (birth control pills). These contain a synthetic estrogen combined with synthetic progesterone, which are taken each day for 3 weeks after the last day of the menstrual period. Placebo pills are taken for the fourth week, to cause a fall in the blood levels of estrogen and progesterone so that menstruation can occur. The birth control pills immediately produce high blood levels of estrogen and progesterone, mimicking the luteal phase and causing negative feedback inhibition of FSH and LH. Thus, no follicles grow and ovulate (so fertilization is prevented), and no corpus luteum can be formed. The newer contraceptive pills have other benefits: they may reduce the risk of endometrial and ovarian cancer, as well as osteoporosis. However, they may also increase the risk of breast cancer, and possibly cervical cancer. Each woman should consult with a physician to weigh the potential benefits and risks in light of her own medical situation and family history.

Review

Male reproductive anatomy & physiology

- male sexual structures
- physiology of an erection
- reproductive problems (ED, BPH)

Gametogenesis


- spermatogenesis
- oogenesis


Steroidogenesis


- Hypothalamic-pituitary-gonadal axis
- negative feedback inhibition of steroidogenesis


23


23















Uterine Fibroids = benign (noncancerous) growths of myometrium, which often appear during childbearing years. ~80% women aged 50 yrs and up

Also called **leiomyomas** (lie-o-my-O-muhs) or **myomas**.

Are NOT associated with an increased risk of uterine cancer (almost never develop into cancer)



UTERINE FIBROIDS

Symptoms:

May have none Heavy menstrual bleeding Pelvic pressure / pain Backache Frequent/difficult urination







CLINICAL APPLICATIONS

The majority of *hysterectomies* (surgical removal of the uterus) are performed because of **uterine fibroids** (**leiomyomas**). These are nonmalignant (noncancerous) neoplasms (growths) in the uterus that also include abundant extracellular matrix. Fibroids can be as small as 10 mm or as large as 20 cm, and produce such symptoms as pelvic discomfort and profuse menstrual bleeding. Uterine fibroids have receptor proteins for estradiol and progesterone, which can stimulate their growth. Because most fibroids are located within the uterine wall, they usually can be surgically removed only by a hysterectomy.

31




Danger of an Out-of-place pregnancy = only uterus & its strong ligaments can support weight of growing fetus. Only endometrium capable of forming a fully functional placenta. All other tissues not compatible for pregnancy.



An ectopic pregnancy is NEVER viable for the embryo AND is lifethreatening for the mother. However, protections for emergency life-saving care for women has been rescinded (<u>June, 2025</u>) due to political issues, putting women's lives in jeopardy across the country.

33



HPV Vaccine - 2006

- Gardasil marketed by Merck & Cervarix by GlaxoSmithKline
- Both are set of 3 vaccinations given over a 6 month period.




Only Gardasil is:


- Effective against 4 strains HPV 2 which cause cancer & 2 which cause warts
- Tested & recommended for 9-26 yr old girls AND boys (younger is better - before sexual exposure!)
- Can get up to 21-26 yrs but protection goes down w/sexual exposure.

https://www.cdc.gov/hpv/parents/vaccine.html

35


HPV infections targeted by vaccine have decreased since vaccination was introduced. Among teen girls! Among young women! HEALTH CARE PROVIDERS should recommend HPV vaccination for all patients at age 11 or 12.* HPV can cause some cancers in women and men HPV vaccination is cancer prevention 1-19 Yacatana stoleard 2.798 1-19 Yaca





IF no fertilization:

- > Corpus luteum breaks down and stops progesterone secretion @day 28.
- ➤ Without **progesterone**, endometrium secretes **prostaglandin**, which cause uterine contractions to expel menstrual tissue.
- Menstrual flow egg and lining shed





- Embryo makes hcg within 1 week (the hormone pregnancy tests detect)
- hCG "rescues" corpus luteum it keeps making progesterone ~ 1month (until placenta forms and takes over progesterone production).


39


CLINICAL APPLICATIONS

Because hCG is secreted by the cells of the chorionic membrane of the embryo, and not by the mother's endocrine glands, all pregnancy tests assay (test) for hCG in urine or blood. Modern pregnancy tests detect the beta subunit of hCG (one of two different polypeptide chains that comprise the protein), which is unique to hCG and provides the least amount of cross-reaction with related hormones. Pregnancy tests use *monoclonal antibodies* (produced by lymphocyte clones; see chapter 11), which are specific for the beta subunit of hCG and are produced by animals such as rabbits injected with hCG. Home pregnancy tests, using monoclonal antibodies that react with hCG in urine, are generally accurate in the week following the first missed menstrual period.









Ovarian cancer (Click HERE to read more)

↑ risk factors include:

- > Genetics (close female relative had it)
- > Have had more ovulations in life (never been on hormonal birth control or been pregnant)
- >Have mutation in the BRCA gene
- > Polycystic ovarian syndrome
- > hormonal problems

↓ risk factors include:

- > not have genetics •
- > no mutation in BRCA gene •
- > fewer ovulations in life •

(have on hormonal birth control - 50% decreased risk

have been pregnant – lower risk)

Question: Why do you think having been on birth control lowers risk of ovarian cancer??

45

Ovarian and Breast Cancer and the BRCA Gene:

BRCA Gene = tumor suppressor gene that normally suppresses tumor growth (a good thing!)

Mutation in BRCA Gene – means the gene does not suppress tumors. Mutation in this gene associated with increased risk for ovarian & breast cancer. People positive for the gene mutation can pass it on to their children. Each child has 50% chance of inheriting the gene mutation.



Ovarian cyst

Angelina Jolie BRCA 1 mutation

Can get blood test for it. BRCA gene mutation test for genetics

CA-125 test = cancer antigen 125 (a non-genetic test)

increased levels of this in blood associated with ↑ risk of ovarian cancer (separate from BRCA gene)

Cancer and the B Increased risk of gene mutation *			
Cancer Type	No mutation	BRCA 1 mutation	BRCA 2 mutation
Female Breast**	12%	65%	57%
Male Breast	<1%	2%	6%
Ovarian***	10%	46%	23%
Prostate ⊀	10%	26%	61%
Pancreatic	1.7%	3%	7%

^{*} Children of parents with BRCA gene mutation have 50% chance of inheriting.

Click HERE for a YouTube video I made about my breast cancer story AND the BRCA gene mutation.



breast cancer double mastectomy part 1

^{* *} Women with BRCA gene mutation have 40% chance of recurrence of cancer if they did not get estrogen blocker treatment or have ovaries removed.

^{* * *} Women with BRCA gene mutation are 100 times more likely to develop cancer in fallopian tubes, before ovaries.

Review

Female reproductive anatomy & physiology

- reproductive structures
- ectopic pregnancy & endometriosis, uterine fibroids
- HPV, warts, cervical cancer, HPV vaccine, ovarian & breast cancer
- review of oogeneis
- menstrual & ovarian cycle
- role of hCG in rescuing corpus luteum in pregnancy
- polycystic ovarian syndrome
- tests for cancer (CA-125 antigen and BRCA gene mutation)

49